



OR-5

CYANO AND CF₃ DERIVATIVES OF 2-ARYL- AND 2-THIENYLQUINAZOLINE: SYNTHESIS AND STUDY OF OPTICAL PROPERTIEST. N. Moshkina,^{1*} Ju. V. Permyakova,¹ E. V. Nosova,^{1,2} G. N. Lipunova,² V. N. Charushin^{1,2}¹ Chemical Technology Institute, UrFU, 620002 Ekaterinburg, 19 Mira St.² Institute of Organic Synthesis, 620219 Ekaterinburg, 22 S. Kovalevskoy St.

*E-mail: tan.moshckina@yandex.ru

Abstract. Organic nonlinear optical (NLO) materials have attracted widespread attention owing to their potential applications in modulation of optical signals, medicine, imaging, laser technology, data storage, telecommunication etc.^{1,2}



The novel 2-thienylquinazoline derivatives **1a-c** containing cyano-group at position 4 as well as quinazolines with 4-cyanophenyl **2a-c** or 4-CF₃-phenyl **2d-f** substituents at position 2 have been synthesized in good yields using early described methods.³⁻⁵


1: Ar = 4-(diethylamino)phenyl (**a**),
4-(diphenylamino)phenyl (**b**),
4-(9H-carbazol-9-yl)phenyl (**c**);

2: R = CN, Ar = 4-(diethylamino)phenyl (**a**),
4-(diphenylamino)phenyl (**b**),
4-(9H-carbazol-9-yl)phenyl (**c**);

R = CF₃, Ar = 4-(dimethylamino)phenyl (**d**),
4-(diphenylamino)phenyl (**e**), **1b** **1c** **2a** **2b** **2c** **2d** **2e** **2f**

4-(9H-carbazol-9-yl)phenyl (**f**).

	φ, %		
	toluene	CHCl ₃	CH ₃ CN
1c	22	10	8
2a	63	63	<1
2b	70	72	16
2c	31	64	20
2d	69	76	12
2e	65	75	31
2f	15	53	28



Emission
solvatochromism

The photophysical properties as well as cyclic voltametry measurements for the quinazoline derivatives **1,2** were studied.

References

- Liu, B., Zhang, H.-L., Liu, J., et al. (2007). Novel pyrimidine-based amphiphilic molecules: synthesis, spectroscopic properties and applications in two-photon fluorescence microscopic imaging. *J. Mater. Chem.* Vol. 17, pp. 2921–2929.
- Cinteza, L.-O., Marinescu, M. (2018). Synthesis and nonlinear studies on selected organic compounds in nanostructured thin films, Advanced surface engineering research, *Intech Open*, doi:10.5772/intechopen.79522.
- Nosova, E. V., Moshkina, T. N., Lipunova, G. N., et al. (2018). Synthesis and photophysical studies of novel 2-[5-(4-diethylaminophenyl)thiophen-2-yl]quinazoline derivatives, *Mendeleev Commun.*, Vol. 28, Iss. 1, pp. 14–16.
- Nosova, E. V., Moshkina, T. N., Lipunova, G. N., et al. (2018). Synthesis and photophysical studies of novel 4-aryl substituted 2-phenyl-, 2-(fluoren-2-yl)- and 2-cymantrenylquinazolines. *Mendeleev Commun.* Vol. 28, Iss. 1, pp. 33–35.
- Achelle, S., RodríguezLópez, J., Guen F.R. (2014). Synthesis and photophysical studies of a series of quinazoline chromophores. *J. Org. Chem.* Vol. 79, pp. 7564–7571.

This work was supported by the Russian Foundation for Basic Research (grant # 18-03-00112) and Russian Science Foundation (project # 19-73-10144)